

Grid diagrams and Khovanov homology

Jean-Marie Droz and Emmanuel Wagner

Abstract

We explain how to compute the Jones polynomial of a link from one of its grid diagrams and we observe a connection between Bigelow's homological definition of the Jones polynomial and Kauffman's definition of the Jones Polynomial. Consequently, we prove that the Maslov grading on the Seidel-Smith symplectic link invariant coincides with the difference between the homological grading on Khovanov homology and the Jones grading on Khovanov homology. We give some evidences for the truth of the Seidel-Smith conjecture.

Introduction

Using symplectic geometry, Seidel and Smith constructed an invariant of oriented links in S^3 [10]. The Seidel-Smith invariant of an oriented link L is defined as the homology $Kh_{symp}^*(L)$ of a chain complex associated to L . The homological grading of this chain complex is denoted by P . They also conjectured that this invariant is isomorphic to the Khovanov link homology $Kh^{*,*}(L)$:

Conjecture (Seidel and Smith). *For all $k \in \mathbb{Z}$,*

$$Kh_{symp}^k(L) \cong \bigoplus_{\substack{(i,j) \in \mathbb{Z}^2 \\ i-j=k}} Kh^{i,j}(L),$$

where i is called the Khovanov homological grading and j the quantum grading.

Manolescu showed [5] that the generators of the Seidel-Smith chain complex are in one-to-one correspondence with the intersection points between homology representative arising in Bigelow's construction of the Jones polynomial [1]. Moreover, this correspondence allows us to endow the Seidel-Smith generators with Bigelow's Jones grading J . Supporting Seidel-Smith conjecture, it has been observed by Manolescu that there is, on small examples, enough generators in the Bigelow construction to have a complex generated by them with homology the Khovanov homology (taking into account the gradings).

The purpose of the present article is to introduce a differential on the graded free abelian group generated by the Bigelow intersection points, also called Bigelow's generators. This aim is achieved by proving that there is an injection of Bigelow's generators into enhanced Kauffman states [4, 11]. In other words, we will see the Seidel-Smith generators as a subset of the generators of the Khovanov chain complex. Moreover, we prove that, as expected, the gradings verify $P = i - j$ and $j = J$. Our main theorem is:

Theorem 1. *There exists a differential δ on the free abelian group B generated by Bigelow's intersection points, that respects J , increases P by 1 and such that the chain complex (B, δ) has homology the Khovanov homology.*

Our differential on the free graded abelian group generated by Bigelow’s generators gives a complex homotopic to the combinatorially defined complex used to compute Khovanov homology [4]. This gives strong supporting evidence for the Seidel-Smith conjecture by generalizing Manolescu’s observation. The main result remains true for odd Khovanov homology [8], see Theorem 8.

The techniques used below are of intrinsic interest. Namely, we develop a combinatorial description of the Jones polynomial and of Khovanov homology in terms of *rectangular diagrams* (see Section 1). In particular, our result gives an alternative proof of the equivalence of Bigelow’s definition of the Jones polynomial. In addition, grid diagrams appear in the combinatorial description of link Floer homology [6]. This new description will be used in future work to investigate the relation between Khovanov type homologies and Heegaard-Floer type homologies (see e.g. [7, 9, 3]).

Outline. In the first section, we introduce all definitions and notations necessary to compute the Jones polynomial from a grid diagram. In the second section, we construct an injection of Bigelow’s generators into enhanced Kauffman’s states. In the third section, we prove relations between the gradings and in the last section we prove our main theorem.

Acknowledgements. We wish to thank Anna Beliakova for interesting discussions and pointing to us the article of Manolescu. The present article would not exist without the kind encouragements of Benjamin Audoux.

1 Definitions and notations

1.1 Grid diagrams and links

A *grid diagram* of size $n \in \mathbb{N} - \{0, 1\}$ is a $(n \times n)$ -grid whose squares may be decorated by either an O or an X so that each column and each row contains exactly one O and one X . The number n is called the *complexity* of the grid diagram. Following [6], we denote by \mathcal{O} the set of O ’s and \mathcal{X} the set of X ’s. The X ’s and the O ’s are called the *punctures* of the grid diagram.

From any grid diagram, one can construct an oriented link diagram. For this purpose, one should join the X and the O in each column by a vertical segment and the O to the X in each row by an horizontal segment that passes under all the vertical segments. We choose the orientation to be from the O ’s to the X ’s on the horizontal lines and from the X ’s to the O ’s on the vertical lines. This produces a *planar rectangular diagram* for an oriented link in S^3 . Any oriented link in S^3 admits a planar rectangular diagram [2]. An example is shown in Figure 1.

1.2 The set \mathcal{G} of the Bigelow’s intersection points

Let D be a planar rectangular diagram of complexity n . For each vertical segment let us draw an oriented curve which winds clockwise around the puncture at the top of the segment and counterclockwise around the puncture at the bottom of the segment. The oriented curves obtained are called *figure-eights*. The figure-eights are very narrow and as short as possible. We assume that the figure-eights intersect transversally and at most twice the horizontal segments, do not intersect each other and have only one transversal self-intersection. We denote by \mathcal{G} the set of unordered n -tuples of




Figure 1: Grid diagram and planar rectangular diagram for the trefoil

intersection points between horizontal segments and (vertical) figure-eights, such that each (vertical) figure-eight and each (horizontal) segment contains exactly one point. We denote by \mathcal{Z} the set of intersection points between figure-eights and horizontal segments. We define $\mathbf{x} \in \mathcal{G}$ ($\mathbf{o} \in \mathcal{G}$, respectively) as the set of points of \mathcal{Z} that are nearest to the X 's (the O 's, respectively), see Figure 2 for an example. To each element $g \in \mathcal{G}$, one can associate a unique n -tuple, $\bar{g} = (\bar{g}_1, \dots, \bar{g}_n)$ in which \bar{g}_i ($i = 1, \dots, n$) is the X , O or crossing nearest to g_i .




Figure 2: Oriented figure-eights and particular elements \mathbf{x} and \mathbf{o} of \mathcal{G}

1.3 Gradings on \mathcal{G}

Three gradings on \mathcal{G} will be defined. We define $\bar{P} : \mathcal{G} \rightarrow \mathbb{Z}$ as follows. Figure-eights are oriented as in Figure 2. Each $z \in \mathcal{Z}$ is given an integer $p(z)$: +1 if the part of the figure-eight on which z sits is oriented upward, 0 otherwise. Given $g = (z_1, \dots, z_n) \in \mathcal{G}$, $\bar{P}(g) = p(z_1) + \dots + p(z_n)$. For example, in Figure 2, $\bar{P}(\mathbf{x}) = 2$ and $\bar{P}(\mathbf{o}) = 2$.

Given finite sets of points A and B in the real plane, we define $\mathcal{I}(A, B)$ to be the number of pairs $(a_1, a_2) \in A$ and $(b_1, b_2) \in B$ such that $a_1 < b_1$ and $a_2 < b_2$. The grading $T : \mathcal{G} \rightarrow \mathbb{Z}$ is defined by $T(g) = \mathcal{I}(g, g)$, for $g \in \mathcal{G}$.

We define a relative grading Q on \mathcal{G} . Consider two elements $g = (g_1, \dots, g_n)$ and $h = (h_1, \dots, h_n)$ in \mathcal{G} . To define the difference $Q(g) - Q(h)$, we consider the loop $\gamma(g, h)$ in the configuration space of n points in \mathbb{R}^2 defined as follows. We start at g , go along the horizontal segments to h , then go back along the vertical figure-eights to g . We can also see $\gamma(g, h)$ as a family of closed immersed oriented curves in \mathbb{R}^2 . Then $Q(g) - Q(h)$ is defined to be the sum of the winding numbers of these closed immersed

curves around the X 's and the O 's. In other words, for each X and O , we count algebraically the number of times each immersed curve goes around the puncture: $+1$ for each time a curve goes around a puncture counterclockwise and -1 for each time a curve goes around a puncture clockwise and we take the sum over all curves and all punctures as relative grading. We define the absolute grading Q by setting $Q(\mathbf{x}) = 0$.

1.4 Normalization of the gradings and the Jones polynomial

We introduce two classical quantities associated to an oriented link diagram D (and hence, to an oriented planar rectangular diagram). Given an oriented link diagram D , we resolve all the crossings of D as in Figure 3, we obtain a disjoint union of oriented




Figure 3: Oriented resolution

circles embedded in \mathbb{R}^2 . We call these circles the *Seifert circles* of D . The *rotation number* of D , denoted by $\text{rot}(D)$, is the sum of the contributions of the circles. The contribution of a Seifert circle is $+1$ if it is oriented counterclockwise and -1 otherwise. Given a crossing c of an oriented link diagram D , we define $w(c)$ as in Figure 4. We define the *writhe* $w(D)$ of D ,

$$w(D) = \sum_{c \text{ crossings of } D} w(c).$$

We denote by n_+ the number of positive crossings and by n_- the number of negative crossings of D . We have $w(D) = n_+ - n_-$.



Negative crossing c , $w(c) = -1$



Positive crossing c , $w(c) = +1$

Figure 4: Crossings

We define the Jones grading J and the homological grading P . Given a planar rectangular diagram D , for an element $g \in \mathcal{G}$ we set

$$J(g) = 2(T(g) - Q(g)) - 2T(\mathbf{x}) + \text{rot}(D) + w(D) \text{ and}$$

$$P(g) = \overline{P}(g) - \overline{P}(\mathbf{x}) - \text{rot}(D) - w(D).$$

We prove below that the Jones polynomial of an oriented planar rectangular diagram D can be written as:

$$V(D)(q) = \sum_{g \in \mathcal{G}} (-1)^{P(g)} q^{J(g)}. \quad (1)$$




Figure 5: From a rectangular diagram to a braid closure

2 Bigelow’s generators and enhanced Kauffman states

2.1 From plat closures to rectangular diagrams

Equation (1) is a simple reformulation of Bigelow’s homological definition of the Jones polynomial. In [1], Bigelow computes the Jones polynomial of a link represented as the plat closure of a braid. Our set \mathcal{G} is seen in the setting of Bigelow’s definition as the set of intersection points between a cycle in homology represented by the figure-eights and another cycle represented by the horizontal segments.

In [5] (Section 3, page 15 and 16), Manolescu explains how the plat closure of a braid can be given as a flattened braid diagram. A rectangular diagram can easily be transformed into a flattened braid diagram. Starting with a planar rectangular diagram in the plane with figure-eights drawn, we apply a diffeomorphism of the plane sending all horizontal segments of the rectangular diagram to consecutive non-intersecting segments on a line, see Figure 5. We obtain a flattened braid diagram. Our definitions of the different gradings are obtained by “pulling back” the gradings originally defined by Bigelow along the diffeomorphism defined above. Notice that we use the notations and normalizations of Manolescu.

2.2 The injection ϕ from Bigelow’s generators to enhanced Kauffman states

Fix an oriented rectangular diagram D , we construct a bijection between the set \mathcal{G} and a subset of the enhanced Kauffman states of an oriented link diagram. Given an unoriented link diagram D , an enhanced Kauffman state is a choice of one resolution for each crossing of D (see Figure 6), together with a choice of orientation on every resulting circle, see Figure 7 for an example. We call a choice of resolution for each crossing of D a *resolution* of D . We define \mathcal{K} to be the set of enhanced Kauffman states.




Figure 6: Kauffman resolutions

Consider an oriented planar rectangular diagram D , we define \mathcal{H} to be the set of enhanced Kauffman states associated to the underlying unoriented link diagram such




Figure 7: Example of an enhanced Kauffman state

that around each crossing, the arcs coming from the resolutions are oriented as in one of the configurations depicted in Figure 8.



Figure 8: Allowed orientations around a resolution in an enhanced Kauffman state

Notice that the enhanced Kauffman states that are not in \mathcal{H} are those for which at least near one crossing the orientation is as in Figure 9.



Figure 9: Forbidden orientations around a resolution in an enhanced Kauffman state

Let D be an oriented planar rectangular diagram, let $g = (g_1, \dots, g_n) \in \mathcal{G}$. We explain how to associate to g an element $\phi(g) \in \mathcal{H}$, see e.g. Figure 11. Thinking of a planar rectangular diagram D as a set of horizontal and vertical segments, we can subdivide all the segments in D at the crossings of D to obtain a set of smaller segments that we denote by $\text{seg}(D)$. To an element $g \in \mathcal{G}$ corresponds an orientation of the segments of $\text{seg}(D)$ by the following two rules:

- A vertical segment s is oriented upward when the intersection point of \bar{g} nearest to the line containing s is lower than s . It is oriented downward otherwise.
- An horizontal segment s is oriented leftward when the intersection point of \bar{g} nearest to the line containing s is at the left of s . It is oriented rightward otherwise.

This means that around an intersection point of \bar{g} the orientation of $\text{seg}(D)$ looks like in Figure 10. See the Figure 11, the diagram in the middle for an another example.

We obtain the enhanced Kauffman state $\phi(g)$ from the orientation on $\text{seg}(D)$ by specifying a resolution of each crossing of D . The orientation on the circles of $\phi(g)$ being induced in the obvious way by the orientation on $\text{seg}(D)$. For a crossing c , if there is no intersection point of g near c , we resolve c in the only way that is coherent with the orientation on $\text{seg}(D)$ (see Figure 3). If there is an intersection point \bar{g}_i of \bar{g} in c , then the resolution depends on the position of g_i . There are four possible cases




Figure 10: Orientation near a \bar{g}_i




Figure 11: An element of \mathcal{G} , the orientation it induces on $\text{seg}(D)$ and its corresponding enhanced Kauffman state $\phi(g)$

and Figure 12 describes how to resolve in each case. Since the orientation on $\text{seg}(D)$ is coherent with our choice of resolutions, the function $g \rightarrow \phi(g)$ is well defined.




Figure 12: Resolution near a g_i

Theorem 2. *Let D be an oriented planar rectangular diagram. The application $\phi : g \mapsto \phi(g)$ defines a bijection between \mathcal{G} and \mathcal{H} .*

Proof. We define a map ψ from \mathcal{H} to \mathcal{G} , see Figure 13 for an example. Given an enhanced Kauffman state h in \mathcal{H} , consider the orientation induced by h on $\text{seg}(D)$. We claim that on a vertical segment of D , there is at most one switch of orientation of the kind shown in Figure 14. This follows from the fact that if there were two, then between them, there would be one switch of the type depicted in Figure 15, which is impossible since enhanced Kauffman state in \mathcal{H} do not contain resolutions oriented as in Figure 9. A similar argument proves that, on each horizontal segment, there is at most one switch of orientation (of the allowed type). It follows that one can associate to an element $h \in \mathcal{H}$, a n -tuple $\psi(h)$ of X 's, O 's or crossings, such that for any $g \in \mathcal{G}$,




Figure 13: Orientation induced on D by an enhanced Kauffman state and associated element of \mathcal{G}



Figure 14: Switch of orientation




Figure 15: Forbidden switch of orientation

$\bar{g} = \overline{\psi(h)}$ implies that g and h induce the same orientation on $\text{seg}(D)$. By using Figure 12, we replace crossings in $\overline{\psi(h)}$ by elements of \mathcal{Z} according to the orientations of the circles around the crossing and replace punctures in $\overline{\psi(h)}$ by their nearest point in \mathcal{Z} . We obtain a n -tuple $\psi(h) \in \mathcal{G}$. Since clearly ϕ and ψ are inverses, ϕ defines a bijection between \mathcal{G} and \mathcal{H} . \square

3 Gradings

3.1 Khovanov homological grading and quantum grading

We introduce two gradings i and j on \mathcal{H} . By the previous theorem, they induce two gradings on \mathcal{G} . Given an element $h \in \mathcal{H}$, consider the underlying resolution $r(h)$ and define $\bar{i}(h)$ to be the number of resolutions in $r(h)$ of the type depicted in Figure 16. Then $i(h) = \bar{i}(h) - n_-$. We define $j(h) = \text{rot}(h) + \bar{i}(h) + n_+ - 2n_-$. Notice that $J(\mathbf{x}) = j(\mathbf{x})$. As a consequence, we define for any $g \in \mathcal{G}$,

$$j(g) = j(\phi(g)) = \text{rot}(\phi(g)) + \bar{i}(\phi(g)) + n_+ - 2n_-,$$

$$i(g) = i(\phi(g)) = \bar{i}(\phi(g)) - n_-.$$

We express the grading $j(g)$ directly from g described as a set of intersection points. More precisely, we decompose the grading j as the sum of three contributions j_1 , j_2 and j_3 .




Figure 16: Resolution used in the definition of Khovanov homological grading on \mathcal{H}

We define $j_1(g)$ to be an algebraic count on the corners of the rectangular diagram. Each corner of the rectangular diagram has a contribution of $+1$ or -1 . Given g , $j_1(g)$ is the sum over all corners of these contributions. For the contributions of each corners, see Figure 17 and Figure 18.



Figure 17: Corners with a contribution to j_1 of $+1$



Figure 18: Corners with a contribution to j_1 of -1

We define $j_2(g)$ to be an algebraic count on the crossings that are near an intersection point of g . In other words, if $g = (g_1, \dots, g_n)$, consider $\bar{g} = (\bar{g}_1, \dots, \bar{g}_n)$. If \bar{g}_i is on a crossing, it has a contribution of $+1$ or -1 depending on the positions of g_i and of the self-intersection of the corresponding figure-eight, see Figure 19 and Figure 20.




Figure 19: Crossings with a contribution to j_2 of $+1$

We define $j_3(g)$ to be an algebraic count on the crossings that do not have a point of g nearby. In other words, if $g = (g_1, \dots, g_n)$, consider $\bar{g} = (\bar{g}_1, \dots, \bar{g}_n)$ and consider the crossings where there is no \bar{g}_i . Each of these crossings has a contribution of $+1$ or -1 depending on the position of the g_i that is on the same figure-eight and on the position of the g_j that is on the same horizontal segment, see Figure 21 and Figure




Figure 20: Crossings with a contribution to j_2 of -1

22. Notice that j_3 does not depend on the position of the self-intersection.




Figure 21: Crossings with a contribution to j_3 of $+1$




Figure 22: Crossings with a contribution to j_3 of -1

From previous definitions, we deduce that, for all $g \in \mathcal{G}$,

$$\begin{aligned} \text{rot}(\phi(g)) &= \frac{j_1(g)}{4} + \frac{j_2(g)}{2} \text{ and} \\ i(g) = i(\phi(g)) &= \frac{j_2(g)}{2} + \frac{j_3(g)}{2} + \frac{n_+ - n_-}{2}. \end{aligned} \quad (2)$$

Hence,

$$j(g) = \frac{j_1(g)}{4} + j_2(g) + \frac{j_3(g)}{2} + \frac{3}{2}(n_+ - n_-), \text{ for all } g \in \mathcal{G}. \quad (3)$$

3.2 Relations between gradings

In this subsection, we prove that the gradings verify $P = i - j$ and $J = j$.

Theorem 3. *Let D be an oriented planar rectangular diagram. For all $g \in \mathcal{G}$,*

$$P(g) = i(g) - j(g).$$

Proof. We denote by \mathcal{G}' the set of unordered n -tuples of intersection points between horizontal segments and vertical figure-eights such that each (vertical) figure-eight contains exactly one point. Observe that \mathcal{G} is a subset of \mathcal{G}' . Moreover, the gradings j_1, j_2 and P on \mathcal{G} extend in a natural way to \mathcal{G}' . Since

$$j(\mathbf{x}) = \text{rot}(D) + w(D), \quad i(\mathbf{x}) = 0 \quad \text{and} \quad P(\mathbf{x}) = -\text{rot}(D) - w(D),$$

we obtain

$$P(\mathbf{x}) = i(\mathbf{x}) - j(\mathbf{x}).$$




Figure 23: Elementary moves

We prove that P and $i - j$ are equal as relative gradings. Notice that one can go from an element of \mathcal{G}' to another by a sequence of elementary moves changing the position of just one intersection point on one figure-eight. It remains to check that P and $i - j$ change by the same amount when such an elementary move is performed. The different cases of elementary moves are presented in Figure 23. Moreover, combining Equations (2) and (3), we have

$$j(g) = \frac{j_1(g)}{4} + \frac{j_2(g)}{2} + i(g) + w(D), \text{ for all } g \in \mathcal{G}.$$

Thus, we need to prove that $\frac{j_1}{4} + \frac{j_2}{2}$ and $-P$ vary by the same amount when an elementary move is performed. This can be checked directly from the pictures in Figure 23. Notice that there are a priori 16 other cases to check corresponding to changes of positions of the self-intersections of the figure-eights in the twelve first elementary moves depicted in Figure 23, but these cases follow from the following observation: A change of position of the self-intersection of the figure-eight together with a change of position of the generator by switching side if it sits between the old and the new self-intersections changes neither $\frac{j_1}{4} + \frac{j_2}{2}$ nor $-P$, see Figure 24.



Figure 24: Change of position of a self-intersection point of a figure-eight

□

Theorem 4. *Let D be an oriented planar rectangular diagram, for all $g \in \mathcal{G}$,*

$$J(g) = j(g).$$

Proof. Since

$$j(\mathbf{x}) = \text{rot}(D) + w(D) = J(\mathbf{x}),$$

it is sufficient to prove that j and J are equal as relative gradings.




Figure 25: The two parts of the figure-eight

For A and B two sets of intersection points, we define $\mathcal{I}^-(A, B)$ as the number of pairs $(a_1, a_2) \in A$ and $(b_1, b_2) \in B$ such that $a_1 < b_1$ and $a_2 > b_2$. Given $g = (g_1, \dots, g_n) \in \mathcal{G}$, we define $Q_{far}(g)$ as follows:

$$Q_{far}(g) = \mathcal{I}(g, \mathbb{X} \cup \mathbb{O}) - \mathcal{I}^-(g, \mathbb{X} \cup \mathbb{O}).$$

We divide any figure-eight in a positive and a negative part by cutting it at its lowest and highest points, Figure 25. We define $Q_{loc}(g)$ as the number of intersection points of g on the positive side of the figure-eight (in blue) minus the number of points on the negative side of the figure-eight (in red). We observe that

$$2P =_{rel} Q_{loc}. \quad (4)$$

(The notation $=_{rel}$ means that the two sides of the equation are equal up to an additive constant.)

The relative grading Q between two generators g and h is defined as the total winding number of a set of closed paths around the punctures. This winding number of a path can be computed by counting intersection points of the path with the horizontal and vertical lines originating from all punctures (with some signs) and dividing by 4. Using this alternative definition of the winding number and examining separately the contribution of the pieces of the path on horizontal lines and on figure-eights, we check that $4Q(g) - 4Q(h) = 2Q_{loc}(g) - 2Q_{loc}(h) + Q_{far}(g) - Q_{far}(h)$, in other words, that, as relative gradings, $4Q =_{rel} 2Q_{loc} + Q_{far}$.

Let us call *good* a line either vertical or horizontal that meets two punctures. We call intersection points of good lines *good points*.

Claim. *We have the equality of relative gradings:*

$$4T - j_3 + \frac{j_1}{2} =_{rel} Q_{far}. \quad (5)$$

Proof. For each pair of one horizontal good line and one vertical good line, we examine the contributions of all pairs of one puncture or intersection point on the vertical line and one puncture or intersection point on the horizontal line to $4T$, j_3 , $\frac{j_1}{2}$ and Q_{far} . We consider different cases according to where the two good lines intersect. We check that for each pair of good lines, the total contribution to Equation (5) from points on the two lines are independent of the positions of the intersection points of g . For example, if the two good lines intersect on a crossing of the rectangular diagram, only $4T$ and j_3 are affected and the contribution of the intersection points to $4T$ and j_3 cancel. Since each puncture or intersection point appears once on a vertical good line and once on an horizontal good line, taking the sum over all possible pairs of good lines gives twice Equation 5. \square

Using Equation (5) divided by two and the definition of j by Equation (3), we get

$$j =_{rel} \frac{j_1}{4} + j_2 + 2T - \frac{Q_{far}}{2} + \frac{j_1}{4}.$$

Using Equation (4) and the formulation of P in terms of j_1 and j_2 , we have

$$-Q_{loc} =_{rel} \frac{j_1}{2} + j_2.$$

Combining the last two equations gives

$$j =_{rel} 2T - \frac{Q_{far}}{2} - Q_{loc}.$$

Since $\frac{Q_{far}}{2} + Q_{loc} =_{rel} 2Q$ and $J =_{rel} 2(T - Q)$, we deduce $j =_{rel} J$ from which $j = J$ follows. \square

4 Relation with Khovanov homology

In [5], Manolescu noticed that, in the case of the trefoil, a free abelian group whose generators are labelled by Bigelow generators and are graded according to the grading of their label, has ranks in the different gradings compatible with having as homology the Khovanov homology. We prove that this observation holds for any link. For this purpose, we start with the Khovanov chain complex, i.e the chain complex whose generators are labelled by enhanced Kauffman states \mathcal{K} and eliminate all generators lying in $\mathcal{K} \setminus \mathcal{H}$. We end up with a chain complex homotopic to the original one, with a set of generators in one-to-one correspondance with \mathcal{G} . This reduction can be done canonically over \mathbb{Q} . Over \mathbb{Z} the reduction seems to depend of some arbitrary choices.

Our reduction to a smaller complex is *canonical* in the following sense. Two oriented rectangular diagrams that are sent to each other by diffeomorphisms of the plane that send vertical segments to vertical segments have isomorphic chain complexes. This implies that the chain complex generated by Bigelow's homological intersection points is computable from the ambient isomorphism type of the flattened braid diagram and is as such a candidate for a geometric interpretation.

Proof. [of Theorem 1] We review a few facts about Khovanov homology (for precise definitions and more [4, 11]). The Khovanov chain complex is a bigraded complex

$$C_{Kh} = \bigoplus_{i,j} C_{Kh}^{i,j}$$

with generators associated to enhanced Kauffman states

$$C_{Kh}^{i,j} = \bigoplus_{s \in \mathcal{K}, i(s)=i, j(s)=j} \mathbb{Z} \cdot s.$$

Observe that $C_{Kh}^i = \bigoplus_{j \in \mathbb{Z}} C_{Kh}^{i,j}$ is generated by all enhanced Kauffman states obtained by orienting the circles in a resolution r such that $i(r) = i$. Hence, as usual, C_{Kh}^i can be seen as $\bigoplus_{r, i(r)=i} V^{\otimes k(r)} \{i + n_+ - n_-\}$ where V is the two dimensional graded \mathbb{Z} -module spanned by 1 and x with $j(1) = 1$ and $j(x) = -1$, $k(r)$ is the number of circles in the resolution r and $\{\cdot\}$ is the shift operator in homological grading. One can identify an element in $V(r) = V^{\otimes k(r)}$ with a choice of orientations of the $k(r)$ circles. A circle oriented counterclockwise corresponds to a 1 and a circle oriented clockwise corresponds to an x . Given a diagram D , with k crossings there are 2^k resolutions of D . One can see these resolutions as lying on the vertices of an hypercube of dimension k . Hence, the vector spaces $V(r)$ are indexed by the vertices of the hypercube. Similarly, each enhanced Kauffman state sits on a vertex of the hypercube. Two enhanced Kauffman states are connected by the differential of Khovanov homology if and only if they are on both ends of an edge of the hypercube and look around a crossing like one of the 18 pairs in Figure 26. Locally, all possible non-zero differentials are depicted in Figure 26. Naturally, one would have to introduce signs to fully specify the differential [4, 11].

We define a new grading R on the Khovanov chain complex as follows. Given an oriented link diagram $D \in \mathbb{R}^2$, consider the underlying oriented 4-valent graph Γ in which each crossing of D is replaced by a 4-valent vertex. Choose a point in each connected component of $\mathbb{R}^2 \setminus \Gamma$. This produces a family of points $(x_i)_{i \in I}$, where I is a finite set. For an enhanced Kauffman state s , we define $R(s)$ to be the winding number of the oriented circles of s around the x_i 's. In Figure 26, the blue cross and the green cross correspond to x_i 's.

The six differentials in the first column of Figure 26 are connecting generators belonging to $\mathcal{K} \setminus \mathcal{H}$. These differentials respect the grading R . In addition, the remaining twelve other types of differentials strictly decrease R . Therefore, the increasing filtration associated to the grading R is respected by the differential of the Khovanov complex. As a consequence, the set of elements of $\mathcal{K} \setminus \mathcal{H}$ together with the part of the differential that respects the grading R is a chain complex. Moreover, this chain complex is homotopic to zero. This follows from the fact that this chain complex is a shifted direct sum of hypercube chain complexes, where each hypercube chain complex is obtained by the usual procedure of flattening an hypercube [7, p.17] and replacing every vertex by a copy of \mathbb{Z} and every arrow by $+\text{Id}$ or $-\text{Id}$. Such an hypercube chain complex is clearly homotopic to zero.




Figure 26: Non-zero differentials in Khovanov complex

We state a lemma in homological algebra generalizing Gaussian elimination which is proved in the Appendix.

Lemma 5. *Given a chain complex $C = \bigoplus_{i \in \mathbb{Z}} C^i$, $C \xrightarrow{\delta} C$, δ of degree $+1$ such that $C = A \oplus B$ as graded abelian group and $\delta = \begin{pmatrix} a & d \\ c & b \end{pmatrix}$ with $A \xrightarrow{a} A$, $B \xrightarrow{d} A$, $A \xrightarrow{c} B$ and $B \xrightarrow{b} B$. Suppose that (B, b) is a chain complex homotopic to zero. Then, for $h : C \rightarrow C$ of degree -1 such that $-\text{Id} = hb + bh$, the chain complex (C, δ) is homotopic to $(A, a + dhc)$.*

Theorem 1 follows from Theorem 3, Theorem 4 and Lemma 5. \square

Corollary 6 (Bigelow [1]). *Given an oriented rectangular diagram D ,*

$$V(D)(q) = \sum_{g \in \mathcal{G}} (-1)^{P(g)} q^{J(g)}$$

is the Jones polynomial.

Corollary 7. *There exists a differential δ on the \mathbb{Q} -vector space B generated by Bigelow's intersection points \mathcal{G} , that respects J , increases P by 1 and such that the chain complex (B, δ) has homology the Khovanov homology. Moreover, over \mathbb{Q} , this differential is canonical.*

Proof. The first part of the statement follows from Theorem 1. The second part follows from the fact that, over \mathbb{Q} , there is a canonical homotopy for Lemma 5. We construct this homotopy. The set of elements of $\mathcal{K} \setminus \mathcal{H}$ together with the part of the differential that leaves the grading R invariant is a chain complex denoted by C . Moreover, this chain complex is a shifted direct sum of hypercube chain complexes, where each hypercube chain complex is obtained by the usual procedure of flattening an hypercube and in this case replacing every vertex by a copy of \mathbb{Q} and every arrow by $+\text{Id}$ or $-\text{Id}$. As explained before, those hypercubes are null homotopic. In particular, given a decomposition of an hypercube of dimension n into two hypercubes of dimension $n - 1$, one can take as homotopy minus the inverse of the differentials between the two hypercubes of dimension $n - 1$. Since there is n ways to split an hypercube of dimension n into two hypercubes of dimension $n - 1$, there are n homotopies of the type described above. We denote them by h_1, h_2, \dots, h_n . Each of them comes from a choice of splitting. We consider $h = \frac{1}{n} \sum_{i=1}^n h_i$; it is a homotopy to zero. Summing over all hypercubes composing the chain complex C , the average homotopies described above, one obtains a homotopy to zero for the whole complex C . Hence, over \mathbb{Q} , there is a canonical choice of homotopy for the application of Lemma 5 and therefore, a canonically defined differential on the set of Bigelow's generators. \square

Theorem 1 and Corollary 7 remain true for the odd Khovanov homology [8]. More precisely, the proof only depends on the fact that one can endow enhanced Kauffman's states with a differential that respects the increasing filtration associated to the grading R and such that the part of the differential that respects the grading R is connecting elements in $\mathcal{K} \setminus \mathcal{H}$. The differential constructed in [8] is up to signs the original Khovanov differential. Hence, we also have the following theorem:

Theorem 8. *There exists a differential δ on the free abelian group B generated by Bigelow's intersection points \mathcal{G} , that respects J , increases P by 1 and such that the chain complex (B, δ) has homology the odd Khovanov homology. Moreover, over \mathbb{Q} , this differential is canonical.*

Appendix

We give the proof of Lemma 5.

Proof. We begin by proving that $(A, a + dhc)$ is a chain complex. Notice that $\delta^2 = 0$ and $b^2 = 0$ imply

$$cd = 0. \tag{6}$$

Moreover, $\delta^2 = 0$ also implies

$$a^2 + dc = 0, \tag{7}$$

$$ad + db = 0, \tag{8}$$

$$ca + bc = 0. \tag{9}$$

Hence,

$$\begin{aligned}
(a + dhc)(a + dhc) &= a^2 + dhca + adhc + dhcdhc \\
&= a^2 + dhca + adhc \\
&= -dc - dhbc - dbhc \\
&= -d(\text{Id} + hb + bh)c \\
&= 0.
\end{aligned}$$

We define a chain map f from (C, δ) to $(A, a + dhc)$ by the formula

$$f = \begin{pmatrix} 1 & dh \end{pmatrix}$$

and a chain map g from $(A, a + dhc)$ to (C, δ) by the formula

$$g = \begin{pmatrix} 1 \\ hc \end{pmatrix}.$$

It is straightforward to check that $f\delta = (a + dhc)f$ and $\delta g = g(a + dhc)$ using (6), (9), (8) and $-\text{Id} = hb + bh$. Moreover, we define a homotopy $H = \begin{pmatrix} 0 & 0 \\ 0 & h \end{pmatrix}$ going from C to C . It follows from (6) and $-\text{Id} = hb + bh$ that $gf - \text{Id} = H\delta + \delta H$. We define H' as follows: $H' = dh^3c$. Let us check that $gf - \text{Id} = H'(a + dhc) + (a + dhc)H'$. We have

$$gf - \text{Id} = \text{Id} + dh^2c - \text{Id} = dh^2c \tag{10}$$

Moreover, we have

$$\begin{aligned}
dh^2c &= -dhbh^2c - dh^2bhc \\
&= 2dh^2c + dbh^3c + dh^3bc \\
&= 2dh^2c - adh^3c - dh^3ca,
\end{aligned}$$

from which we deduce,

$$dh^2c = dh^3ca + adh^3c. \tag{11}$$

Hence combining (10) and (11), we have

$$gf - \text{Id} = dh^3ca + adh^3c.$$

Moreover,

$$\begin{aligned}
H'(a + dhc) + (a + dhc)H' &= dh^3c(a + dhc) + (a + dhc)dh^3c \\
&= dh^3ca + adh^3c.
\end{aligned}$$

□

References

- [1] S. Bigelow, *A homological definition of the Jones polynomial*, Invariants of knots and 3-manifolds (Kyoto, 2001), 29–41 (electronic), Geom. Topol. Monogr., 4, Geom. Topol. Publ., Coventry, 2002.
- [2] I. A. Dynnikov, *Arc-presentations of links. Monotonic simplification*, math.GT/0208153v3

- [3] E. Grigsby and S. Wehrli, *On the Colored Jones Polynomial, Sutured Floer homology and Knot Floer homology*, math.GT/0807.1432
- [4] M. Khovanov, *A categorification of the Jones polynomial*, Duke Math. J. 101-3 (2000) 359–426.
- [5] C. Manolescu, *Nilpotent slices, Hilbert schemes and the Jones polynomial*, Duke Math. J. 132 (2006), no. 2, 311–369.
- [6] C. Manolescu, P. Ozsvath, Z. Szabo, D. Thurston, *On combinatorial link Floer homology*, math.GT/0610559
- [7] P. Ozsváth and Z. Szabó, *On the Heegaard Floer homology of branched double-covers*, Adv. Math. 194 (2005), no. 1, 1–33.
- [8] P. Ozsváth, J. Rasmussen, Z. Szabó, *Odd Khovanov homology*, math.GT/0710.4300
- [9] J. Rasmussen, *Knot polynomials and knot homologies*, Geometry and topology of manifolds, 261–280, Fields Inst. Commun., 47, Amer. Math. Soc., Providence, RI, 2005.
- [10] P. Seidel and I. Smith, *A link invariant from the symplectic geometry of nilpotent slices*, Duke Math. J. 134 (2006), no. 3, 453–514.
- [11] O. Viro, *Remarks on definition of Khovanov homology* math.GT/0202199v1

Jean-Marie Droz, *Institut für Mathematik, Universität Zürich, Winterthurerstrasse 190, CH-8057 Zürich, Switzerland.*
 E-mail address: jdroz@math.unizh.ch

Emmanuel Wagner, *Department of Mathematics, University of Aarhus, DK-8000, Denmark.*
 E-mail address: wagner@imf.au.dk